

RESEARCH ARTICLE

Open Access

# Differential disease resistance response in the barley necrotic mutant *nec1*

Anete Keisa, Krista Kanberga-Silina, Ilva Nakurte, Laura Kunga and Nils Rostoks\*

## Abstract

**Background:** Although ion fluxes are considered to be an integral part of signal transduction during responses to pathogens, only a few ion channels are known to participate in the plant response to infection. CNGC4 is a disease resistance-related cyclic nucleotide-gated ion channel. *Arabidopsis thaliana* CNGC4 mutants *hlm1* and *dnd2* display an impaired hypersensitive response (HR), retarded growth, a constitutively active salicylic acid (SA)-mediated pathogenesis-related response and elevated resistance against bacterial pathogens. Barley CNGC4 shares 67% aa identity with AtCNGC4. The barley mutant *nec1* comprising of a frame-shift mutation of CNGC4 displays a necrotic phenotype and constitutively over-expresses *PR-1*, yet it is not known what effect the *nec1* mutation has on barley resistance against different types of pathogens.

**Results:** *nec1* mutant accumulated high amount of SA and hydrogen peroxide compared to parental cv. Parkland. Experiments investigating *nec1* disease resistance demonstrated positive effect of *nec1* mutation on non-host resistance against *Pseudomonas syringae* pv. *tomato* (*Pst*) at high inoculum density, whereas at normal *Pst* inoculum concentration *nec1* resistance did not differ from wt. In contrast to augmented *P. syringae* resistance, penetration resistance against biotrophic fungus *Blumeria graminis* f. sp. *hordei* (*Bgh*), the causal agent of powdery mildew, was not altered in *nec1*. The *nec1* mutant significantly over-expressed race non-specific *Bgh* resistance-related genes *BI-1* and *MLO*. Induction of *BI-1* and *MLO* suggested putative involvement of *nec1* in race non-specific *Bgh* resistance, therefore the effect of *nec1* on *mlo-5*-mediated *Bgh* resistance was assessed. The *nec1/mlo-5* double mutant was as resistant to *Bgh* as *Nec1/mlo-5* plants, suggesting that *nec1* did not impair *mlo-5* race non-specific *Bgh* resistance.

**Conclusions:** Together, the results suggest that *nec1* mutation alters activation of systemic acquired resistance-related physiological markers and non-host resistance in barley, while not changing rapid localized response during compatible interaction with host pathogen. Increased resistance of *nec1* against non-host pathogen *Pst* suggests that *nec1* mutation may affect certain aspects of barley disease resistance, while it remains to be determined, if the effect on disease resistance is a direct response to changes in SA signaling.

## Background

To date, numerous lesion mimic mutants (LMM) have been characterized in *Arabidopsis thaliana*, rice and maize [1,2]. Frequently, LMM display enhanced disease resistance, constitutive expression of pathogenesis-related responses and an altered hypersensitive response (HR). Molecular mechanisms triggering the onset of cell death underlying the lesions mimic phenotype might have common features with HR-associated cell death observed during pathogen infection [3]. Although a direct link between HR and plant disease resistance is

often questioned [3,4], it is evident that LMM can clarify numerous aspects of plant-pathogen interactions at the molecular level.

Although several barley mutants with necrotic leaf spots have been reported [5], only very few LMM phenotypes of barley have been traced down to a particular gene. The best known examples of barley LMM are *mlo* [6,7], and the recently characterized *necS1* (*HvCAX1*) [8], which apart from displaying a necrotic phenotype also shows enhanced disease resistance against fungal pathogens. The barley mutant *nec1* comprising of a mutated cyclic nucleotide gated ion channel 4 (CNGC4) exhibits the necrotic phenotype and over-expresses the pathogenesis-related gene *PR-1* [9]. *A. thaliana* CNGC4

\* Correspondence: nils.rostoks@lu.lv  
Faculty of Biology, University of Latvia, 4 Kronvalda Boulevard, Riga, LV-1586, Latvia

mutants *dnd2* and *hlm1* which are orthologous to barley *nec1* mutants display enhanced resistance to virulent bacterial pathogens [10,11]. HvCNGC4 shares 67% aa identity with AtCNGC4 [9], suggesting that a similarly to *dnd2* in *A. thaliana* *nec1* mutation may affect barley disease resistance.

Bacterial diseases of barley have been described, although the mechanisms of resistance have not been studied in detail [12,13]. Apparently, there is no race-specific resistance to bacterial pathogens: thus, only PAMP-triggered immunity is operational, even though cultivar-dependent differences in infection rates have been reported for bacterial kernel blight caused by *Pseudomonas syringae* [14]. Significant over-production of salicylic acid (SA) upon *P. syringae* infection in barley suggests that barley resistance to non-host bacterial pathogens is achieved through a SA-mediated defense pathway [15].

Bacterial pathogens of *Arabidopsis* are commonly used as a model system for plant-pathogen interaction studies. However, fungal pathogens are the causal agents of economically more deleterious and widespread diseases in barley. Powdery mildew is caused by the biotrophic fungus *Blumeria graminis* f. sp. *hordei* (*Bgh*). This is among the best studied barley diseases, and extensive details are available on both the race specific or race non-specific powdery mildew resistance mechanisms [16]. Race non-specific resistance of barley to *Bgh* is a cell wall-based resistance forbidding fungal penetration into a host cell [17]. Penetration resistance is triggered by the ROR2 protein, presumably directing secretion vesicle trafficking to the fungal penetration site [18]. Race non-specific penetration resistance is fully attained only in the absence of the trans-membrane protein MLO which is a negative regulator of ROR2 [19]. Functional MLO protein employs  $Ca^{2+}$  and CaM signaling to ensure fungal penetration into host cells. Mutations negatively affecting MLO binding with CaM render barley more resistant against *Bgh* [19,20], while overexpression of another trans-membrane protein, BI-1, counteract *mlo*-triggered *Bgh* resistance in a  $Ca^{2+}$  and CaM signaling-dependent manner [21,22]. Although the interdependence of  $Ca^{2+}$ /CaM signaling and race non-specific *Bgh* resistance in barley is well established, so far no  $Ca^{2+}$  permeable ion-channel has been shown to participate in *Bgh* resistance or susceptibility.

Race specific resistance of barley against *Bgh* requires the presence of plant *R*-genes called *Ml* genes. In contrast to race non-specific *Bgh* resistance, race specific resistance usually permits fungal penetration into the host cell, but restricts further spread of the fungus by triggering plant cell death [16]. Both types of powdery mildew resistance have been shown to incorporate reactive oxygen species (ROS) signaling elements, such as

increased accumulation of  $H_2O_2$  and/or superoxide [23,24].  $H_2O_2$  acts as a principal signaling molecule initiating cell death during incompatible race-specific barley-*Bgh* interaction [24]. Early accumulation of  $H_2O_2$  in mesophyll cells underlying attacked epidermal cells is proposed to be critical for the establishment of race specific resistance [25,26]. In race non-specific interactions,  $H_2O_2$  plays a distinct role from that observed for HR induction. In *mlo*-triggered resistance,  $H_2O_2$  most likely ensures host cell wall fortification, thus preventing fungal penetration [23,27].

In this study, disease resistance of barley LMM *nec1* mutants displaying necrotic leaf spots was analyzed. Although *NEC1* has been shown to encode cyclic nucleotide gated ion channel 4 (CNGC4) and to over-express the defense-related *PR-1* gene [9], the effect of *nec1* mutation on barley disease resistance has not yet been characterized. This study shows that *nec1* mutation triggers the induction of  $H_2O_2$  and SA, restricts *Bgh* microcolony formation and affects non-host resistance against *Pseudomonas syringae* applied at high inoculum density, whereas it has no effect on *Bgh* penetration efficiency or *mlo*-dependent race non-specific *Bgh* resistance.

## Results

### *nec1* mutant exhibits constitutive activation of $H_2O_2$ and salicylic acid

The *nec1* allele in cultivar Parkland was initially described as a natural mutation [28], which was confirmed by identification of a MITE insertion in an intron of the *NEC1* gene that caused alternative splicing and a predicted non-functional protein [9]. The *nec1* mutant line GSHO 1284 and a parental variety Parkland were genotyped with DArT markers [29]. Only 2.2% of 1131 DArT loci were polymorphic, suggesting that the mutant is essentially isogenic to Parkland (data not shown). All described experiments were performed with Parkland and its mutant *nec1* accession GSHO 1284.

As it was found that *nec1* significantly over-expressed pathogenesis related genes [9], it was investigated whether *nec1* plants spontaneously display also other SAR-related signals such as altered accumulation of reactive oxygen species and over-accumulation of SA. Spectrofluorimetric analysis of whole-leaf extracts of two week old *nec1* plants with a fully developed lesion mimic phenotype and the parental line Parkland showed a three-fold higher overall level of  $H_2O_2$  in the mutant (data not shown).

To ascertain whether the elevated overall amount of  $H_2O_2$  in *nec1* plants affected  $H_2O_2$  accumulation during *Bgh* infection, overall  $H_2O_2$  amount in *nec1* and wt plants was assessed at 12 h and 36 h after inoculation with a virulent mixed population of *Bgh*. The analysis


did not reveal considerable changes in the H<sub>2</sub>O<sub>2</sub> content of wt plants during the first 36 h after inoculation, whereas *nec1* mutants showed a slight, statistically non-significant increase in H<sub>2</sub>O<sub>2</sub> levels at 36 h after inoculation (Figure 1).

H<sub>2</sub>O<sub>2</sub> accumulation and *PR-1* expression is known to be associated with SA-dependent signaling. Therefore, the SA content of *nec1* and wt plants was also measured. HPLC assay confirmed that levels of free SA and conjugated SA were four- and fifteen-fold higher, respectively, in *nec1* than in wt plants (Figure 2).


#### Resistance of the *nec1* mutant to *Pseudomonas syringae*

Barley resistance to the non-host bacterial pathogen *Pseudomonas syringae* likely employs SA-mediated defense pathway [15]. Therefore, the constitutive activation of SA signaling in *nec1* might contribute to its non-host resistance. *nec1* plants were inoculated with *P. syringae* pv. *tomato* (*Pst*) at two inoculum densities -  $8 \times 10^4$  and  $6 \times 10^7$  cfu ml<sup>-1</sup> using vacuum infiltration technique. At day 3 after infiltration with  $6 \times 10^7$  cfu ml<sup>-1</sup> of *Pst* the amount of bacteria in *nec1* was reduced, whereas Parkland had accumulated ca. 6-fold higher amount of *Pst* making the difference in bacterial growth between wt and *nec1* statistically highly significant ( $p = 0.01$ , Student's *t*-test) at this stage of infection (Figure 3A). Inoculation with *Pst* at lower inoculum density ( $8 \times 10^4$  cfu ml<sup>-1</sup>) did not reveal any differences in resistance between *nec1* and wt plants (Figure 3A).

Ion leakage measurements were also performed to characterize the effect of *Pst* infection on *nec1* and



**Figure 1** Time course of whole leaf H<sub>2</sub>O<sub>2</sub> accumulation in *nec1* and wt plants after *Bgh* infection. *nec1* mutation triggers H<sub>2</sub>O<sub>2</sub> over-accumulation in barley in the absence of pathogen infection, but it does not alter time course of H<sub>2</sub>O<sub>2</sub> production in response to *Bgh* infection. Error bars represent the standard deviation of means ( $n = 5$  per data point).




**Figure 2** Level of free and conjugated SA in *nec1* and wt plants. *nec1* mutant contains significantly higher level of conjugated, as well as free SA compared to parental cv. Parkland. SA content was analyzed using reverse-phase high performance liquid chromatography in leaf tissue extracts of 14 day old plants. Average values from three biological replicates are presented, each consisting of three technical replicates. Error bars represent standard deviation.

Parkland. Vacuum infiltration with *Pst* at lower inoculum density ( $8 \times 10^4$  cfu ml<sup>-1</sup>) did not elicit cell death in either *nec1* or Parkland (Figure 3B). In contrast to inoculation with lower *Pst* density, inoculation with *Pst* at  $6 \times 10^7$  cfu ml<sup>-1</sup> elicited differential response in *nec1* and wt. Tissue samples from *nec1* plants inoculated with *Pst* at  $6 \times 10^7$  cfu ml<sup>-1</sup> displayed more pronounced ion leakage suggesting an increased cell death in *nec1* after infection (Figure 3B).

#### Resistance of *nec1* mutant to powdery mildew *Blumeria graminis* f.sp. *hordei*


Since *nec1* plants exhibited constitutively active defense responses, the role of *nec1* in basal resistance against *Bgh* was assessed. Due to their basal resistance, even susceptible barley cultivars are able to restrict infection to some extent. In order to assess the effect of *nec1* mutation on basal *Bgh* resistance, microcolony formation was examined. *nec1* supported formation of significantly ( $p < 0.001$ , *t*-test) smaller number of *Bgh* colonies compared to wt plants (Figure 4). To further test, if restricted formation of *Bgh* microcolonies on *nec1* derived from the rapid and effective localized response precluding fungal penetration or from post-invasive defense impeding further fungal development, we examined *nec1* *Bgh* penetration resistance. The effect of *nec1* mutation on *Bgh* penetration resistance was characterized as the proportion of interaction sites that had formed *Bgh* haustoria to the total number of *Bgh* spores



**Figure 3 Response of *nec1* to non-host pathogen *Pseudomonas syringae* pv. *tomato* applied at low and high inoculum densities.** Panel A. Growth of *Pseudomonas syringae* pv. *tomato* in *nec1* and parental cv. Parkland was monitored immediately and 3 days after vacuum infiltration with *Pst* applied at inoculum densities of  $8 \times 10^4$  or  $6 \times 10^7$  cfu ml<sup>-1</sup>. For mock inoculation plants were infiltrated with 10 mM MgCl<sub>2</sub>. Infection was expressed as number of colony forming units (cfu) per gram of fresh leaves (FW). Due to the high between-experiment variation, results of one representative experiment out of four independent experiments are shown. Error bars represent standard deviation. At high inoculum density ( $6 \times 10^7$  cfu ml<sup>-1</sup>) bacterial cfu number in *nec1* at the day 3 was significantly ( $p < 0.01$ , Student's *t*-test) lower than in wt. Panel B. Progression of cell death in *nec1* and Parkland after infection with *Pseudomonas syringae* pv. *tomato* in the experiment shown in panel A. *nec1* mutation showed increased electrolyte leakage in barley inoculated with non-host bacteria *Pst* at  $6 \times 10^7$  cfu ml<sup>-1</sup>. Measurements of electrolyte leakage were taken every two hours during 24 hour period and at 48 hours after inoculation. Error bars represent standard deviation.


that had germinated at 48 hpi. *nec1* plants permitted almost identical entry and haustoria establishment rate of *Bgh* as the parental line (71% and 74% *Bgh* penetration efficiency respectively,  $p = 0.64$ , Student's *t*-test).

Basal *Bgh* resistance has been shown to be tightly linked to the molecular mechanisms of race-specific *Bgh*



**Figure 4 *Bgh* microcolony formation on *nec1* and wt plants.** Excised segments of barley leaves were inoculated with a virulent *Bgh* isolate. Microcolony formation was inspected microscopically 4 days post infection and infection rate was expressed as a number of microcolonies per cm<sup>2</sup> leaf area. Figure reflects data from two independent experiments. Error bars represent standard deviation. Infection frequency significantly differs between *nec1* and Parkland ( $p < 0.001$ , *t*-test).


resistance triggered by different *Mla* alleles [30,31]. *HvRbohA* and *HvRacB* are known to participate in basal as well as race-specific *Bgh* resistance [32-34]. The expression of these genes was characterized using real-time quantitative PCR. Relative mRNA abundance of the analyzed genes was not affected by *nec1* mutation (Figure 5) indirectly suggesting that *nec1* may be independent from effector-triggered immunity that ensure rapid localized *Bgh* resistance.




**Figure 5 Effect of *nec1* mutation on expression of powdery mildew resistance related genes.** Transcript abundance of powdery mildew resistance related genes in *nec1* mutants was determined by quantitative real time PCR. mRNA abundance of *HvMLO* and *HvBI-1* is significantly increased in *nec1*. Error bars represent standard deviation.

***nec1* mutation alters expression of *BI-1* and *MLO*, but does not affect *mlo-5*-triggered race non-specific powdery mildew resistance**

Different powdery mildew resistance types employ at least partially distinctive molecular pathways: thus, a particular gene can have a significant role in one *Bgh* resistance strategy, while having only a marginal or no effect on another *Bgh* resistance type [35]. To find out, if *nec1* mutation affected *mlo*-triggered race non-specific *Bgh* resistance, the expression of *MLO* and *BI-1* genes was analyzed using real-time quantitative PCR. Loss of functional *MLO* protein renders barley almost fully resistant against *Bgh*, whereas *BI-1* over-expression in *mlo* mutants leads to restoration of susceptibility against *Bgh* [22] and, in fact, *BI-1* is required for full susceptibility of barley to powdery mildew [36]. Furthermore, over-expression of *MLO* in wild type plants leads to super susceptibility against *Bgh* [20]. Significant over-expression of both *MLO* and *BI-1* in *nec1* plants was observed (Figure 5). To further test whether *nec1* mutation had any effect on race non-specific powdery mildew resistance conferred by *mlo-5* mutation, *Bgh* penetration resistance of *nec1/mlo-5* double mutant was characterized. Similar to *mlo-5* mutant, *nec1/mlo-5* plants were almost fully resistant to *Bgh*, allowing establishment of fungal haustoria only at less than 2% of interaction sites (Figure 6). In addition, the H<sub>2</sub>O<sub>2</sub> content of whole-leaf extracts from *nec1/mlo-5* double mutants was analyzed. While the *nec1* mutant showed markedly increased accumulation of H<sub>2</sub>O<sub>2</sub> compared to wt *NEC1* plants, the experiment did not reveal a significant effect of *mlo-5* mutation on H<sub>2</sub>O<sub>2</sub> over-accumulation in *nec1* (Figure 7).



**Figure 6 Effects of *nec1* mutation on *mlo-5* triggered *Bgh* penetration resistance.** Fourteen days old plants were inoculated with 10-20 conidia per mm<sup>2</sup> and at 48 h post inoculation infected leaves were harvested and *Bgh* penetration efficiency was assessed. At least 100 interaction sites per variant were observed. Error bars represent standard deviation.



**Figure 7 Effect of *mlo-5* mutation on H<sub>2</sub>O<sub>2</sub> accumulation in barley mutant *nec1*.** *mlo-5* mutation does not affect over-accumulation of H<sub>2</sub>O<sub>2</sub> in *nec1* mutant. H<sub>2</sub>O<sub>2</sub> content was determined spectrofluorimetrically in leaf extracts of wt, *nec1*, *mlo-5* and *nec1/mlo-5* double mutants. Error bars represent standard deviation.

**Discussion**

Despite the fact that ion fluxes are known to play an important role in early signaling events during plant-pathogen interaction [37-39], to date only several plant ion channels have been shown to participate in plant disease resistance or plant-pathogen interaction signal transduction. The cyclic nucleotide gated ion channel (CNGC) gene family is one of the best-represented among the disease resistance-related ion channels. CNGC mutants *dnd1* (AtCNGC2), *dnd2* and *hlm1* (AtCNGC4) and *cpr22* (AtCNGC11/12) exhibit a wide range of pathogen resistance [10,11,40,41]. Mutations affecting AtCNGC4 enhance resistance of *Arabidopsis thaliana* against certain pathotypes of *Pseudomonas syringae* and *Botrytis cinerea* [10,11,42]. Although the effect of CNGC mutations on resistance against bacterial and oomycete pathogens is well-studied in *Arabidopsis*, little is known about the role of these genes in non-host resistance and also about the functions of CNGCs in disease resistance of economically important monocot plant species such as barley. Here we show that similarly to *dnd2* in *A. thaliana* [10], *nec1* in barley activates constitutive over-accumulation of SA. High level of SA contributes to enhanced disease resistance of *dnd2* to virulent *Pseudomonas syringae* pv. *tomato* [10,42] and this resistance requires functional *PAD4* [43], which is one of the central genes in SA-mediated effector-triggered immunity (ETI) [44] and SAR [45]. Although disease resistance pathways seem to be largely conserved among monocots and dicots [46-49], the position of SA in monocot immunity is ambiguous. Some monocots, such as rice, contain high endogenous SA levels [50]

and SA is not required for *PR*-gene induction in rice upon infection [51]. Ineffectiveness of externally applied SA on induction of *PR*-genes has also been observed in barley [15] and wheat [52], however, inoculation with non-host bacteria *Pseudomonas syringae* triggers SA accumulation in barley [15]. Taking into account that such differences occur in the SA mediated resistance signaling among monocots and dicots, it is interesting to see whether mutation affecting SA mediated disease resistance in *A. thaliana* is also involved in barley disease resistance. The present study analyzed the effect of the *nec1* (*HvCNGC4*) mutation on barley resistance against *Pseudomonas syringae* pv. *tomato* and *Blumeria graminis* f. sp. *hordei*.

Mutation in the *NEC1* gene affected barley non-host resistance against *Pseudomonas syringae* pv. *tomato*. Bacterial growth in *nec1* plants was delayed at the initial phase of infection, if plants were inoculated with bacteria at high inoculum density. At the same time the increased electrolyte leakage suggested somewhat enhanced cell death, even though the conductivity values were much lower than reported for typical HR. Thus, electrolyte leakage data in *nec1* were generally in agreement with the expected “defense, no death” phenotype characteristic of *hlm1/dnd2* mutants, although differences between *nec1* and *hlm1/dnd2* mutants may exist in this respect. Non-host resistance is predicted to share common defense responses with host resistance - either basal (PAMP-triggered immunity, PTI) or ETI [53,54]. The choice of which layer of immunity is activated upon a particular interaction with non-host pathogen seems to be case specific [55-57]. Therefore molecular mechanisms leading to changes in non-host resistance of *nec1* to *P. syringae* pv. *tomato* might have also had an effect on interaction with host pathogens. This prompted the assessment of the role of *nec1* mutation in resistance to powdery mildew caused by the fungal pathogen *Blumeria graminis* f. sp. *hordei*. *nec1* restricted *Bgh* microcolony formation, while not affecting *Bgh* penetration or *mlo-5* triggered resistance to *Bgh*. Interestingly, despite the fact that *nec1* did not impede *mlo-5* mediated race non specific resistance to *Bgh*, *MLO* and *BI-1* mRNA abundance was significantly increased in barley *nec1* plants (Figure 5). Significant over-expression of *MLO* and *BI-1* might result from general activation of cell death-related signaling pathways and systemic immunity responses rather than from activation of particular powdery mildew resistance. Together these observations suggest that *nec1* mutation most likely affects PTI and non-host resistance related responses and it is not associated with rapid localized defense responses required to prevent fungal penetration.

HR related cell death is suggested to serve in plant immunity as a factor triggering activation of SAR [4,58]. Spontaneous cell death might elicit constitutive activation of SAR related signaling pathway in *nec1*. Previously *nec1* has been shown to constitutively over-express *PR-1a* and  $\beta$ -1,3-glucanase [9] - molecular markers of SAR. This study confirmed the constitutive activation of SA-related signaling pathways in *nec1* mutants, since significant over-accumulation of H<sub>2</sub>O<sub>2</sub> and SA in *nec1* plants was detected. In *Arabidopsis thaliana*, non-host resistance against some types of pathogens involves SA signaling [59-61]. In barley, a substantial increase in SA levels has been observed after infection with *Pseudomonas syringae* pv. *syringae*, but not after inoculation with non-host fungus *Blumeria (Erysiphe) graminis* f. sp. *tritici* [15] or host pathogen *Bgh* [23] suggesting a differential role of SA in barley resistance against different pathogens. Constitutive activation of the SA-related defense pathway may contribute to differential resistance of *nec1* mutant against non-host bacteria *Pst* and virulent host pathogen *Bgh*. However, the cause for SA over-accumulation needs further investigation, and it remains to be determined, if SA-independent pathways are activated in *nec1* mutant similarly to *Arabidopsis hlm1/dnd2* mutant.

## Conclusions

*nec1* mutation increased resistance against the non-host bacterial pathogen *Pseudomonas syringae* pv. *tomato* applied at high inoculum density and it also inhibited microcolony formation of host pathogen *Blumeria graminis* f.sp. *hordei*, but its penetration resistance to *Bgh* or race non-specific *Bgh* resistance pathways were not impaired. The differential disease resistance response of *nec1* plants might result from the activation of specific resistance pathways differentiating between various types of pathogens. SA-dependent signaling pathways have previously been shown to participate in disease resistance against certain types of pathogens, while not affecting others. *nec1* mutant displays constitutive activation of systemic acquired resistance-related signals such as over-accumulation of hydrogen peroxide and SA, as well as over-expression of *PR-1*. It remains to be determined, if constitutive activation of SA related signaling is the main reason for the differential disease resistance of *nec1* mutant.

## Methods

### Plants

Plants for all experiments were grown in an environmental growth chamber at 22°C under long-day (16 h day, 8 h night), medium light (ca. 150  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) conditions. The barley necrotic mutant *nec1* (GSHO

1284) containing a MITE insertion in the gene for *Cyclic Nucleotide Gated Ion Channel 4 (CNGC4)* [9] has previously been described as a natural mutant in cv. Parkland [28]. Both cv. Parkland and GSHO 1284 are completely susceptible to powdery mildew. *mlo-5* and *nec1* double mutant was obtained by crossing accession GSHO 1284 with NGB 9276 carrying the *mlo-5* allele in the cv. Carlsberg II background [62]. Plants homozygous for *nec1* and *mlo-5* alleles were confirmed by genotyping the respective mutations and F<sub>4</sub> plants were used for all experiments. Barley accessions GSHO 1284 and Parkland were obtained from USDA ARS National Small Grains Germplasm Research Facility (Aberdeen, Idaho, USA), and NGB 9276 was obtained from Nordic Genetic Resources Center (Alnarp, Sweden).

#### **Infection with *Pseudomonas syringae* pv. *tomato***

To study *nec1* non-host resistance against *Pseudomonas syringae* pv. *tomato*, leaves of 14 day old *nec1* plants were vacuum infiltrated with a bacterial suspension in 10 mM MgCl<sub>2</sub>. Bacterial suspension was applied at normal concentration  $8 \times 10^4$  and high concentration  $6 \times 10^7$  cfu ml<sup>-1</sup>, since low concentration inoculum typically applied for infection of host plants can have minor or no effect on non-host species [63]. For mock inoculation 10 mM MgCl<sub>2</sub> was used for infiltration. Immediately after infiltration plants were covered with plastic bags to maintain high humidity and kept in dark for 1 h. After an hour plants were transferred to growth conditions described above. Bacterial growth was monitored at day 3 post inoculation by dilution plating of homogenized plant tissue. Leaves were briefly sterilized with 70% ethanol before homogenization. *Pseudomonas syringae* pv. *tomato* was obtained from the German microbial type collection (accession 50315).

#### **Cell death measurements**

Cell death was quantified by electrolyte leakage assay performed as described by Dellagi et al. (1998) with minor modifications [64]. In brief, plants were vacuum infiltrated with *Pst* as described above and incubated in dark at high humidity for an hour. Five mm leaf disks were collected and washed with distilled water for 1 h and then transferred to a tube with 6.5 ml distilled water. Conductivity was measured with conductivity meter handylab LF11 (Schott Instruments). Each sample contained 4 leaf disks from 4 plants and at each data point 4 independent replicates were measured.

#### **Fungal material, inoculation and calculation of penetration efficiency**

Two week old plants of *nec1* and cv. Parkland were inoculated with 10-20 conidia per mm<sup>2</sup> from virulent mixed population of powdery mildew multiplied on cv.

Parkland. For the characterization of penetration efficiency, infected barley leaves were harvested 48 h post inoculation and cleared for 24 h in 98% ethanol. Penetration efficiency was calculated as a ratio of interaction sites with haustoria formation and the total number of spores with developed appressoria. The overall penetration efficiency for the particular barley line was an average from three replicates containing at least 100 interaction sites each.

*Bgh* microcolony formation was examined on 5 cm long leaf middle segments, which were laid flat on 0.5% agar in water (w v<sup>-1</sup>) plates with adaxial surface facing up and were inoculated with mixed population of powdery mildew multiplied on cv. Parkland. Each plate contained leaves from both *nec1* and cv. Parkland plants to compensate for uneven inoculation. *Bgh* microcolonies were microscopically scored 4 days post inoculation. Experiment was repeated twice with 14 independent samples per barley line in each experiment.

#### **H<sub>2</sub>O<sub>2</sub> detection and quantification**

Hydrogen peroxide was quantified spectrofluorometrically [65]. Briefly, 1 g of freshly harvested leaves from two week old barley plants were frozen in liquid nitrogen and ground in 50 mM Hepes-KOH buffer containing 1 mM EDTA and 5 mM MgCl<sub>2</sub> (pH 7.5). After centrifugation for 10 minutes at 13000 g, the supernatant was transferred to a new centrifuge tube and an equal volume of chloroform:methanol (volume ratio 2:1) solution was added. After centrifugation for 3 minutes at 13000 g, the upper aqueous phase was transferred to a new centrifuge tube and 50 mM Hepes-KOH buffer solution (pH 7.5) containing 0.5 mM homovanillic acid and 15 U horseradish peroxidase VI was added to a final volume of 3 ml. Samples were incubated at room temperature for 30 minutes before fluorescence measurements were taken (excitation at 315 nm, emission at 425 nm). Fluorescence was measured with a FluoroMax3 spectrofluorometer (Horiba Scientific, Japan). For quantification of the H<sub>2</sub>O<sub>2</sub> a standard curve with a range of 100 μM - 1 nM was applied. Sample correction for quenching was performed by adding a known sample amount to a 10 nM H<sub>2</sub>O<sub>2</sub> solution.

#### **Quantification of free and conjugated salicylic acid**

The SA content in leaf tissue extracts was analyzed using reverse-phase high performance liquid chromatography (HPLC). Each sample contained leaf tissue from 3 two week old plants. Samples were prepared essentially as described [66]. Briefly, 0.45 g barley leaf tissue was homogenized in liquid nitrogen and sequentially extracted using 90% and 100% methanol. Extraction was repeated twice and two supernatant fractions were pooled and dried. The residue was resuspended in 1 ml of 5% acetic acid. As an internal standard for SA

recovery correction, samples were selectively spiked with 50 µg per g FW 3-hydroxy benzoic acid (3-HBA) [66].

For the quantification of free SA, 1 ml of ethylacetate:cyclopentane:isopropanol (50:50:1) was added. The sample was thoroughly mixed and the upper phase (approximately 1 ml) was transferred to a new 2 ml tube. The aqueous phase was then re-extracted, as described previously, and both organic phases (approximately 2 ml) were pooled. The resulting solution was vacuum-dried and thoroughly resuspended in 0.9 ml of mobile phase. This suspension was filtered through a 0.20 µm filter.

The aqueous phase containing the SAG fraction was acidified with HCl to pH 1.0 and boiled for 30 min to separate free SA from conjugated SA. The released SA was then extracted with the organic mixture and treated as above.

Chromatographic analysis was performed on a modular HPLC system, Agilent 1100 series, consisting of quaternary pump, autosampler, column thermostat and both UV and fluorescence detectors (Agilent Technologies, Germany). Separation was achieved on a Zorbax Eclipse XDB-C18 (Agilent Technologies, Germany) column 4.6 × 250 mm, 5 µm. Column temperature was maintained at 40°C. The mobile phase was prepared by mixing acetonitrile:20 mM NaH<sub>2</sub>PO<sub>4</sub> (pH 3.0 with acetic acid), in a volume ratio 25:75. The mobile phase flow rate was 1.0 ml min<sup>-1</sup>. Injection volume was 100 µl. The UV/VIS detector was set to 237 nm and 303 nm and the fluorescence detector to an excitation wavelength of 297 nm and an emission wavelength of 407 nm. Results were evaluated by a ChemStation Plus (Agilent, Germany).

#### RNA extraction

For RNA extraction, 5 cm long segments of cotyledon leaf from two week old plants of necrotic mutant *nec1* and parental *cv. Parkland* were frozen in liquid nitrogen immediately after harvesting. Total RNA was extracted from frozen leaf tissues using Trizol reagent. Each RNA sample was extracted from a pool of five plants, and three biological replicates of each barley line (15 plants in total) were used for expression analysis of *BI-1*, *MLO*, *HvRACB* and *HvRbohA* genes in *nec1* and *cv. Parkland* plants. Integrity of the extracted RNA was monitored using non-denaturing agarose gel electrophoresis. Quantity of purified total RNA was monitored using spectrophotometer NanoDrop ND-1000 (NanoDrop products, USA). One to two µg of the extracted RNA was treated with DNaseI (Fermentas, Vilnius, Lithuania) following the manufacturer's instructions and afterwards purified using chloroform-ethanol extraction.

#### Reverse transcription and quantitative real-time PCR

cDNA was synthesized with oligo (dT)<sub>18</sub> primers in a total volume of 10 µl containing 1 µg of total RNA

**Table 1 Quantitative real-time PCR primer sequences used in the study**

Primer	Sequence	Reference
HvBI_cw1	CGATGATCTCCTGCGTGTCG	This study *
HvBI_ccw1	TACCTCGGTGGCCTGCTCTC	This study *
HvGAPDH_cw1	CGTTCATCACCCAGCTACTAC	[67]
HvGAPDH_ccw1	CAGCCTTGTCTTGTGTCAGTG	[67]
MLO_F1	GTCGAGCCCAGCAACAAGTTCTTC	This study *
MLO_R1	ACCACCACCTTCATGATGCTCAG	This study *
HvrbohA_F1	CCGATCAGATGTATGCTCCA	[33]
HvrbohA_R1	CAGAAGGCATTGAAGCCAGT	[33]
HvRACB_L01	GGTAGACAAAGAACAAGGCGGAAGT	This study *
HvRACB_R01	CACAAGGCAGGAAGAAGAAATCA	This study *

\* Primers were designed using Primer 3 software [68] using the following gene sequences as a template: *HvBI* (HarvEST21 Unigene 3323; AJ290421); *MLO* (HarvEST21 Unigene 6351; Z83834); *HvRacB* (HarvEST21 Unigene 5202; AJ344223)

using the RevertAid H Minus First Strand cDNA synthesis kit (Fermentas, Vilnius, Lithuania).

For quantitative real-time PCR, aliquots of cDNA were amplified on an ABI Prism 7300 instrument (Applied Biosystems, Foster City, CA, USA) using the Maxima SYBR Green PCR kit (Fermentas, Vilnius, Lithuania) in a total volume of 20 µl containing 2 µl of cDNA and 0.3 µM primers (Table 1). The reaction was carried out as follows: initial denaturing step for 15 min at 95°C followed by 35 cycles of 15 s at 94°C, 30 s at 60°C and 45 s at 72°C (data acquisition step). Standard curves for the quantification of the transcript levels were calculated from serial dilutions of appropriate cDNA fragments amplified from *cv. Parkland*. Transcript levels of the studied genes were expressed as a percentage of *HvGAPDH* transcript value in the same sample. Combined values of two technical replicates of the three biological replicates (n = 6) were used to calculate the average values and standard deviations. Analysis of variance (ANOVA) of transcript abundance between the mutant and the corresponding parent was done in Microsoft Excel (Redmond, WA, USA).

#### Acknowledgements

The study was funded by Latvian Council of Science grant Z-6142-090, European Social Fund project 2009/0224/1DP/1.1.1.2.0/09/APIA/VIAA/055 and University of Latvia grant ZP-59. AK and LK are recipients of the European Social Fund scholarships (projects 2009/0138/1DP/1.1.2.1.2/09/IPIA/VIAA/004 and 2009/0162/1DP/1.1.2.1.1/09/IPIA/VIAA/004, respectively). Authors are grateful to the anonymous reviewers for their suggestions that helped to improve the manuscript.

#### Authors' contributions

AK designed and performed the study and drafted the manuscript. KKS and LK performed the disease resistance tests and gene expression analyses. IN performed HPLC analysis and helped to draft the manuscript. NR designed and performed the study and wrote the final manuscript. All authors have read and approved the submitted manuscript.



Received: 15 September 2010 Accepted: 15 April 2011  
Published: 15 April 2011

## References

- Lorrain S, Vaillieu F, Balague C, Roby D: Lesion mimic mutants: keys for deciphering cell death and defense pathways in plants? *Trends Plant Sci* 2003, **8**:263-271.
- Moeder W, Yoshioka K: Lesion mimic mutants: A classical, yet still fundamental approach to study programmed cell death. *Plant Signal Behav* 2008, **3**:764-767.
- Mur LA, Kenton P, Lloyd AJ, Ougham H, Prats E: The hypersensitive response; the centenary is upon us but how much do we know? *J Exp Bot* 2008, **59**:501-520.
- Heath MC: Hypersensitive response-related death. *Plant Mol Biol* 2000, **44**:321-334.
- Lundqvist U, Franckowiak J, Konishi T: New and revised descriptions of barley genes. *Barley Genetics Newsletter* 1997, **26**:22.
- Buschges R, Hollricher K, Panstruga R, Simons G, Wolter M, Frijters A, Van Daelen R, Van Der Lee T, Diergarde P, Groenendijk J, et al: The barley *Mlo* gene: A novel control element of plant pathogen resistance. *Cell* 1997, **88**:695-705.
- Wolter M, Hollricher K, Salamini F, Schulze Lefert P: The *mlo* resistance alleles to powdery mildew infection in barley trigger a developmentally controlled defence mimic phenotype. *Mol Gen Genet* 1993, **239**:122-128.
- Zhang L, Lavery L, Gill U, Gill K, Steffenson B, Yan G, Chen X, Kleinohfs A: A cation/proton-exchanging protein is a candidate for the barley *NecS1* gene controlling necrosis and enhanced defense response to stem rust. *Theor Appl Genet* 2008, **118**:385-397.
- Rostoks N, Schmierer D, Mudie S, Drader T, Brueggeman R, Caldwell DG, Waugh R, Kleinohfs A: Barley necrotic locus *nec1* encodes the cyclic nucleotide-gated ion channel 4 homologous to the *Arabidopsis HLM1*. *Mol Genet Genomics* 2006, **275**:159-168.
- Jurkowski GI, Smith RK Jr, Yu IC, Ham JH, Sharma SB, Klessig DF, Fengler KA, Bent AF: *Arabidopsis DND2*, a second cyclic nucleotide-gated ion channel gene for which mutation causes the "defense, no death" phenotype. *Mol Plant Microbe Interact* 2004, **17**:511-520.
- Balague C, Lin B, Alcon C, Flottes G, Malmstrom S, Kohler C, Neuhaus G, Pelletier G, Gaymard F, Roby D: HLM1, an essential signaling component in the hypersensitive response, is a member of the cyclic nucleotide-gated channel ion channel family. *Plant Cell* 2003, **15**:365-379.
- Mathre DE: *Compendium of barley diseases* St. Paul, Minnesota: The American Phytopathological Society; 1997.
- Georgakopoulos DG, Sands DC: Epiphytic populations of *Pseudomonas syringae* on barley. *Canadian Journal of Microbiology* 2010, **38**:111-114.
- Martinez-Miller C, Braun SL, Siemsen SL, Sands DC: Etiology of basal kernel blight of barley caused by *Pseudomonas syringae* pv. *syringae*. *Canadian Journal of Plant Pathology* 1997, **19**:337-346.
- Vallelian-Bindschedler L, Metraux JP, Schweizer P: Salicylic acid accumulation in barley is pathogen specific but not required for defense-gene activation. *Molecular Plant-Microbe Interactions* 1998, **11**:702-705.
- Hueckelhoven R, Kogel KH: Reactive oxygen intermediates in plant-microbe interactions: Who is who in powdery mildew resistance? *Planta* 2003, **216**:891-902.
- Schulze-Lefert P: Knocking on the heaven's wall: pathogenesis of and resistance to biotrophic fungi at the cell wall. *Curr Opin Plant Biol* 2004, **7**:377-383.
- Underwood W, Somerville SC: Focal accumulation of defences at sites of fungal pathogen attack. *J Exp Bot* 2008, **59**:3501-3508.
- Bhat RA, Miklis M, Schmelzer E, Schulze-Lefert P, Panstruga R: Recruitment and interaction dynamics of plant penetration resistance components in a plasma membrane microdomain. *Proc Natl Acad Sci USA* 2005, **102**:3135-3140.
- Kim MC, Panstruga R, Elliott C, Mueller J, Devoto A, Yoon HW, Park HC, Cho MJ, Schulze-Lefert P: Calmodulin interacts with MLO protein to regulate defence against mildew in barley. *Nature* 2002, **416**:447-450.
- Ihara-Ohori Y, Nagano M, Muto S, Uchimiyama H, Kawai-Yamada M: Cell death suppressor *Arabidopsis bax inhibitor-1* is associated with calmodulin binding and ion homeostasis. *Plant Physiol* 2007, **143**:650-660.
- Hueckelhoven R, Dechert C, Kogel KH: Overexpression of barley BAX inhibitor 1 induces breakdown of *mlo*-mediated penetration resistance to *Blumeria graminis*. *Proc Natl Acad Sci USA* 2003, **100**:5555.
- Hueckelhoven R, Fodor J, Preis C, Kogel KH: Hypersensitive cell death and papilla formation in barley attacked by the powdery mildew fungus are associated with hydrogen peroxide but not with salicylic acid accumulation. *Plant Physiol* 1999, **119**:1251-1260.
- Hueckelhoven R: Powdery mildew susceptibility and biotrophic infection strategies. *FEMS Microbiol Lett* 2005, **245**:9-17.
- Hueckelhoven R, Fodor J, Trujillo M, Kogel KH: Barley *Mla* and *Rar* mutants compromised in the hypersensitive cell death response against *Blumeria graminis* f.sp. *hordei* are modified in their ability to accumulate reactive oxygen intermediates at sites of fungal invasion. *Planta* 2000, **212**:16-24.
- Vanacker H, Carver TL, Foyer CH: Early H<sub>2</sub>O<sub>2</sub> accumulation in mesophyll cells leads to induction of glutathione during the hyper-sensitive response in the barley-powdery mildew interaction. *Plant Physiol* 2000, **123**:1289-1300.
- Collins NC, Thordal-Christensen H, Lipka V, Bau S, Kombrink E, Qiu JL, Huckelhoven R, Stein M, Freialdenhoven A, Somerville SC, et al: SNARE-protein-mediated disease resistance at the plant cell wall. *Nature* 2003, **425**:973-977.
- Fedak G, Tsuchiya T, Helgason SB: Use of monotelotrisomics for linkage mapping in barley. *Can J Gen Cytol* 1972, **14**:949-957.
- Wenzl P, Carling J, Kudrna D, Jaccoud D, Huttner E, Kleinohfs A, Kilian A: Diversity Arrays Technology (DArT) for whole-genome profiling of barley. *Proc Natl Acad Sci USA* 2004, **101**:9915-9920.
- Caldo RA, Nettleton D, Peng J, Wise RP: Stage-specific suppression of basal defense discriminates barley plants containing fast- and delayed-acting *Mla* powdery mildew resistance alleles. *Mol Plant Microbe Interact* 2006, **19**:939-947.
- Shen QH, Saijo Y, Mauch S, Biskup C, Bieri S, Keller B, Seki H, Ulker B, Somsich IE, Schulze-Lefert P: Nuclear activity of MLA immune receptors links isolate-specific and basal disease-resistance responses. *Science* 2007, **315**:1098-1103.
- Schultheiss H, Dechert C, Kogel KH, Hueckelhoven R: A small GTP-binding host protein is required for entry of powdery mildew fungus into epidermal cells of barley. *Plant Physiol* 2002, **128**:1447-1454.
- Trujillo M, Altschmied L, Schweizer P, Kogel KH, Huckelhoven R: Respiratory burst oxidase homologue A of barley contributes to penetration by the powdery mildew fungus *Blumeria graminis* f. sp. *hordei*. *J Exp Bot* 2006, **57**:3781-3791.
- Wong HL, Pinontoan R, Hayashi K, Tabata R, Yaeno T, Hasegawa K, Kojima C, Yoshioka H, Iba K, Kawasaki T, et al: Regulation of rice NADPH oxidase by binding of Rac GTPase to its N-terminal extension. *Plant Cell* 2007, **19**:4022-4034.
- Peterhansel C, Freialdenhoven A, Kurth J, Kolsch R, Schulze-Lefert P: Interaction analyses of genes required for resistance responses to powdery mildew in barley reveal distinct pathways leading to leaf cell death. *Plant Cell* 1997, **9**:1397-1409.
- Eichmann R, Bischof M, Weis C, Shaw J, Lacomme C, Schweizer P, Duchkov D, Hensel G, Kumlehn J, Huckelhoven R: BAX INHIBITOR-1 is required for full susceptibility of barley to powdery mildew. *Mol Plant Microbe Interact* 2010, **23**:1217-1227.
- McDowell JM, Dangi JL: Signal transduction in the plant immune response. *Trends Biochem Sci* 2000, **25**:79-82.
- Dangi JL, Jones JG: Plant pathogens and integrated defence responses to infection. *Nature* 2001, **411**:826-833.
- Numberger T, Scheel D: Signal transmission in the plant immune response. *Trends Plant Sci* 2001, **6**:372-379.
- Clough SJ, Fengler KA, Yu IC, Lippok B, Smith RK Jr, Bent AF: The *Arabidopsis dnd1* "defense, no death" gene encodes a mutated cyclic nucleotide-gated ion channel. *Proc Natl Acad Sci USA* 2000, **97**:9323-9328.
- Yoshioka K, Moeder W, Kang HG, Kachroo P, Masmoudi K, Berkowitz G, Klessig DF: The chimeric Arabidopsis CYCLIC NUCLEOTIDE-GATED ION CHANNEL11/12 activates multiple pathogen resistance responses. *Plant Cell* 2006, **18**:747-763.
- Genger RK, Jurkowski GI, McDowell JM, Lu H, Jung HW, Greenberg JT, Bent AF: Signaling pathways that regulate the enhanced disease resistance of *Arabidopsis* "defense, no death" mutants. *Mol Plant Microbe Interact* 2008, **21**:1285-1296.

43. Jirage D, Zhou N, Cooper B, Clarke JD, Dong X, Glazebrook J: **Constitutive salicylic acid-dependent signaling in *cpr1* and *cpr6* mutants requires PAD4.** *The Plant Journal* 2001, **26**:395-407.
44. Wiermer M, Feys BJ, Parker JE: **Plant immunity: the EDS1 regulatory node.** *Curr Opin Plant Biol* 2005, **8**:383-389.
45. Durrant WE, Dong X: **Systemic acquired resistance.** *Annu Rev Phytopathol* 2004, **42**:185-209.
46. Chern MS, Fitzgerald HA, Yadav RC, Canlas PE, Dong X, Ronald PC: **Evidence for a disease-resistance pathway in rice similar to the *NPR1*-mediated signaling pathway in *Arabidopsis*.** *Plant Journal* 2001, **27**:101-113.
47. Cooper B, Clarke JD, Budworth P, Kreps J, Hutchison D, Park S, Guimil S, Dunn M, Luginbuhl P, Ellero C, et al: **A network of rice genes associated with stress response and seed development.** *Proc Natl Acad Sci USA* 2003, **100**:4945-4950.
48. Qiu D, Xiao J, Ding X, Xiong M, Cai M, Cao Y, Li X, Xu C, Wang S: **OsWRKY13 mediates rice disease resistance by regulating defense-related genes in salicylate- and jasmonate-dependent signaling.** *Mol Plant Microbe Interact* 2007, **20**:492-499.
49. Humphry M, Bednarek P, Kemmerling B, Koh S, Stein M, Gobel U, Stuber K, Pislewska-Bednarek M, Loraine A, Schulze-Lefert P, et al: **A regulon conserved in monocot and dicot plants defines a functional module in antifungal plant immunity.** *Proc Natl Acad Sci USA* 2010.
50. Chen Z, Iyer S, Caplan A, Klessig DF, Fan B: **Differential accumulation of salicylic acid and salicylic acid-sensitive catalase in different rice tissues.** *Plant Physiol* 1997, **114**:193-201.
51. Yang Y, Qi M, Mei C: **Endogenous salicylic acid protects rice plants from oxidative damage caused by aging as well as biotic and abiotic stress.** *The Plant Journal* 2004, **40**:909-919.
52. Molina A, Gorlach J, Volrath S, Ryals J: **Wheat genes encoding two types of PR-1 proteins are pathogen inducible, but do not respond to activators of systemic acquired resistance.** *Mol Plant Microbe Interact* 1999, **12**:53-58.
53. Mysore KS, Ryu CM: **Nonhost resistance: how much do we know?** *Trends Plant Sci* 2004, **9**:97-104.
54. Jones JD, Dangl JL: **The plant immune system.** *Nature* 2006, **444**:323-329.
55. Lipka U, Fuchs R, Lipka V: ***Arabidopsis* non-host resistance to powdery mildews.** *Curr Opin Plant Biol* 2008, **11**:404-411.
56. Niks RE, Marcel TC: **Nonhost and basal resistance: how to explain specificity?** *New Phytol* 2009, **182**:817-828.
57. Schweizer P: **Nonhost resistance of plants to powdery mildew New opportunities to unravel the mystery.** *Physiological and Molecular Plant Pathology* 2007, **70**:3-7.
58. Kombrink E, Schmelzer E: **The hypersensitive response and its role in local and systemic disease resistance.** *European Journal of Plant Pathology* 2001, **107**:69-78.
59. Yun BW, Atkinson HA, Gaborit C, Greenland A, Read ND, Pallas JA, Loake GJ: **Loss of actin cytoskeletal function and EDS1 activity, in combination, severely compromises non-host resistance in *Arabidopsis* against wheat powdery mildew.** *The Plant Journal* 2003, **34**:768-777.
60. van Wees SC, Glazebrook J: **Loss of non-host resistance of *Arabidopsis NahG* to *Pseudomonas syringae* pv. *phaseolicola* is due to degradation products of salicylic acid.** *The Plant Journal* 2003, **33**:733-742.
61. Zimmerli L, Stein M, Lipka V, Schulze-Lefert P, Somerville S: **Host and non-host pathogens elicit different jasmonate/ethylene responses in *Arabidopsis*.** *The Plant Journal* 2004, **40**:633-646.
62. Jorgensen H: **Discovery, characterization and exploitation of *Mlo* powdery mildew resistance in barley.** *Euphytica* 1992, **63**:141-152.
63. Lopez-Solanilla E, Bronstein PA, Schneider AR, Collmer A: **HopPtoN is a *Pseudomonas syringae* Hrp (type III secretion system) cysteine protease effector that suppresses pathogen-induced necrosis associated with both compatible and incompatible plant interactions.** *Mol Microbiol* 2004, **54**:353-365.
64. Dellagi A, Brisset MN, Paulin JP, Expert D: **Dual role of desferrioxamine in *Erwinia amylovora* pathogenicity.** *Mol Plant Microbe Interact* 1998, **11**:734-742.
65. Jimenez A, Creissen G, Kular B, Firmin J, Robinson S, Verhoeven M, Mullineaux P: **Changes in oxidative processes and components of the antioxidant system during tomato fruit ripening.** *Planta* 2002, **214**:751-758.
66. Aboul-Soud MAM, Cook K, Loake GJ: **Measurement of salicylic acid by a High-Performance Liquid Chromatography procedure based on ion-exchange.** *Chromatographia* 2004, **59**:129-133.
67. Horvath H, Rostoks N, Brueggeman R, Steffenson B, von Wettstein D, Kleinhofs A: **Genetically engineered stem rust resistance in barley using the *Rpg1* gene.** *Proc Natl Acad Sci USA* 2003, **100**:364-369.
68. Untergasser A, Nijveen H, Rao X, Bisseling T, Geurts R, Leunissen JA: **Primer3Plus, an enhanced web interface to Primer3.** *Nucleic Acids Res* 2007, **35**:W71-W74.

doi:10.1186/1471-2229-11-66

**Cite this article as:** Keisa et al.: **Differential disease resistance response in the barley necrotic mutant *nec1*.** *BMC Plant Biology* 2011 **11**:66.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

